

# Intramammary Product Comparison

Professional Veterinarian Services Technical Bulletin



Dry Cow Treatment Options					Indications			
Product/ Manf.	Rx Required	Min. Dry Period Length	Milk	Meat	Staph aureus	Strep agalactiae	Strep dysgalactiae	Strep uberis
<b>Albadry Plus™</b> (novobiocin penicillin) Zoetis		30 Days	72 Hours	30 Days	X	X		
<b>Dry-Clox®</b> (cloxacillin benzathine) BIVI	X	30 Days	0 Hours	30 Days	X	X		
<b>go-dry®</b> (penicillin) Hanford's US Vet		14 Days	72 Hours	14 Days		X		
<b>Orbenin-DC®</b> (benzathine cloxacillin) Merck	X	28 Days	0 Hours	28 Days	X	X		
<b>Quartermaster</b> (penicillin-dihydrostreptomycin) WG Critical Care	X	42 Days	96 Hours	60 Days	X			
<b>Spectramast® DC</b> (ceftiofur hydrochloride) Zoetis	X	30 Days	0 Hours	16 Days	X		X	X
<b>ToMORROW®</b> (cephapirin benzathine) BIVI		30 Days	72 Hours	42 Days	X	X		

Lactating Treatment Options				Indications						Dosage
Product	Rx Required	Milk	Meat	Coagulase-neg. staph	Staph aureus	Strep agalactiae	Strep dysgalactiae	Strep uberis	E. coli	
<b>Amoxi-Mast®</b> (amoxicillin) Merck	X	60 Hours	12 Days		X	X				1 syringe; may repeat every 12 hours for a maximum of 3 treatments
<b>Dariclox®</b> (sodium cloxacillin) Merck	X	48 Hours	10 Days		X	X				1 syringe; may repeat every 12 hours for a maximum of 3 treatments
<b>Pirsue®</b> (pirlimycin hydrochloride) Zoetis	X	36 Hours	9-21* Days		X	X	X	X		1 syringe; may repeat every 24 hours for up to 8 consecutive days
<b>PolyMast™</b> (hetacillin potassium) BIVI	X	72 Hours	10 Days		X	X	X		X	1 syringe; may repeat every 24 hours for a maximum of 3 treatments
<b>Spectramast® LC</b> (ceftiofur hydrochloride) Zoetis	X	72 Hours	2 Days	X			X		X	1 syringe; may repeat every 24 hours for up to 8 consecutive days
<b>ToDAY®</b> (cephapirin sodium) BIVI		96 Hours	4 Days		X	X				1 syringe every 12 hours for a maximum of 2 treatments

\*9-day slaughter withhold required after last treatment when used for 2 days;  
21-day slaughter withhold when treatment period has been 3-8 days.

